

Literatur zum Beitrag

Christoph Leithner, Martin Kenda, Michael Scheel und Thomas Els

Zerebrale Bildgebung nach Herzstillstand und Reanimation

1. Myrhaugen Tjelmeland IB, Alm-Kruse K, Granser JT et al.: Importance of reporting survival as incidence: a cross-sectional comparative analysis on out-of-hospital cardiac arrest registry data from Germany and Norway. *BMJ Open* 2022; 12(2): e058381
2. Cronberg T, Greer DM, Lilja G et al.: Brain injury after cardiac arrest: from prognostication of comatose patients to rehabilitation. *Lancet Neurol* 2020; 19(7): 611–22
3. Velly L, Perlberg V, Boulier T et al.: Use of brain diffusion tensor imaging for the prediction of long-term neurological outcomes in patients after cardiac arrest: a multicenter, international, prospective, observational, cohort study. *Lancet Neurol* 2018; 17(4): 317–26
4. Moseby-Knappe M, Pellis T, Dragancea I: Head computed tomography for prognostication of poor outcome in comatose patients after cardiac arrest and targeted temperature management. *Resuscitation* 2017; 119: 89–94
5. Streitberger KJ, Endisch C, Ploner CJ, et al.: Timing of Brain Computed Tomography and Accuracy of Outcome Prediction after Cardiac Arrest. *Resuscitation* 2019; 145: 8–14
6. Greer DM, Wu O: Neuroimaging in cardiac arrest prognostication. *Semin Neurol* 2017; 37:66–74
7. Bender A: S1-Leitlinie Hypoxisch-ischämische Enzephalopathie im Erwachsenenalter. Deutsche Gesellschaft für Neurologie 2018; Leitlinien für Diagnostik und Therapie in der Neurologie
8. Nolan JP, Sandroni C, Böttiger BW et al.: European Resuscitation Council and European Society of Intensive Care Medicine Guidelines 2021: Post-resuscitation care. *Intensive Care Med* 2021; 47(4): 369–421
9. Inamasu J, Miyatake S, Tomioka H et al.: Subarachnoid hemorrhage as a cause of out-of-hospital cardiac arrest: a prospective computed tomography study. *Resuscitation* 2009; 80(9): 977–80
10. Arnaut M, Mongardon N, Deye N et al.: Out-of-hospital cardiac arrest from brain cause: epidemiology, clinical features, and outcome in a multicenter cohort. *Crit Care Med* 2015; 43(2): 453–60
11. Kenda M, Scheel M, Kemmling A et al.: Automated Assessment of Brain CT After Cardiac Arrest-An Observational Derivation/Validation Cohort Study. *Crit Care Med* 2021; 49(12): e1212-e1222
12. Inamasu J, Miyatake S, Suzuki M et al.: Early CT signs in out-of-hospital cardiac arrest survivors: Temporal profile and prognostic significance. *Resuscitation* 2010; 81(5): 534–8
13. Choi KS, Won YD, Yi HJ et al.: Therapeutic and prognostic implications of subarachnoid hemorrhage in patients who suffered cardiopulmonary arrest and underwent cardiopulmonary resuscitation during an emergency room stay. *Clin Neurol Neurosurg* 2013; 115: 2088–93
14. Scheel M, Storm C, Gentsch A et al.: The prognostic value of gray-white-matter ratio in cardiac arrest patients treated with hypothermia. *Scand J Trauma Resusc Emerg Med* 2013; 21: 23
15. Kirsch K, Heymel S, Günther A et al.: Prognostication of neurologic outcome using gray-white-matter-ratio in comatose patients after cardiac arrest. *BMC Neurol* 2021; 21(1): 456
16. In YN, Lee IH, Park JS et al.: Delayed head CT in out-of-hospital cardiac arrest survivors: Does this improve predictive performance of neurological outcome? *Resuscitation* 2022; 172: 1–8
17. Beekman R, Maciel CB, Ormseth CH et al.: Early head CT in post-cardiac arrest patients: A helpful tool or contributor to a self-fulfilling prophecy? *Resuscitation* 2021; 165: 68–76
18. Caraganis A, Mulder M, Kempainen RR et al.: Interobserver Variability in the recognition of hypoxic-ischemic brain injury on computed tomography soon after out-of-hospital cardiac arrest. *Neurocrit Care* 2020; 33: 414–21
19. Metter RB, Rittenberger JC, Guyette FX et al.: Association between a quantitative CT scan measure of brain edema and outcome after cardiac arrest. *Resuscitation* 2011; 82(9): 1180–5
20. Oh JH, Choi SP, Zhu JH et al.: Differences in the gray-to-white matter ratio according to different computed tomography scanners for outcome prediction in post-cardiac arrest patients receiving target temperature management. *PLoS One* 2021; 16(10): e0258480
21. Hanning U, Bernhard Sporns P, Lebedz P et al.: Automated assessment of early hypoxic brain edema in non-enhanced CT predicts outcome in patients after cardiac arrest. *Resuscitation* 2016; 104: 91–4
22. Friberg H, Cronberg T, Dünser MW et al.: Survey on current practices for neurological prognostication after cardiac arrest. *Resuscitation* 2015; 90: 158–62
23. Lopez Soto C, Dragoi L, Heyn CC et al.: Imaging for neuroprognostication after cardiac arrest: Systematic review and meta-analysis. *Neurocrit Care* 2020; 32: 206–16
24. Lee BK, Jeung KW, Song KH et al.: Prognostic values of gray matter to white matter ratios on early brain computed tomography in adult comatose patients after out-of-hospital cardiac arrest of cardiac etiology. *Resuscitation* 2015; 96: 46–52
25. Youn CS, Callaway CW, Rittenberger JC, et al.: Combination of initial neurologic examination, quantitative brain imaging and electroencephalography to predict outcome after cardiac arrest. *Resuscitation* 2017; 110: 120–5
26. Scarpino M, Lolli F, Lanzo G et al.: ProNeCA study group: Neurophysiology and neuroimaging accurately predict poor neurological outcome within 24 hours after cardiac arrest: The ProNeCA prospective multicentre prognostication study. *Resuscitation* 2019; 143: 115–23
27. Schick A, Prekker ME, Kempainen RR et al.: Association of hypoxic ischemic brain injury on early CT after out of hospital cardiac arrest with neurologic outcome. *Am J Emerg Med* 2022 54: 257–62
28. Els T, Kassubek J, Kubalek R Klisch J: Diffusion-weighted MRI during early global cerebral hypoxia: a predictor for clinical outcome? *Acta Neurol Scand* 2004: 110: 361–7

29. Greer D, Scripko P, Bartscher J et al.: Serial MRI changes in comatose cardiac arrest patients. *Neurocrit Care* 2011; 14: 61–7.
30. Snider SB, Fischer D, McKeown ME, et al. Regional Distribution of Brain Injury After Cardiac Arrest: Clinical and Electrographic Correlates. *Neurology* 2022; 22: 98(12): e1238-e1247
31. Choi SP, Park KN, Park HK et al.: Diffusion-weighted magnetic resonance imaging for predicting the clinical outcome of comatose survivors after cardiac arrest: A cohort study. *Critical Care* 2010; 14(1): R17. doi: 10.1186/cc8874
32. Christophe C, Fonteyne C, Zierysen F et al.: Value of MR imaging of the brain in children with hypoxic coma. *Am J Neuroradiol* 2002; 23: 716–23
33. Takahashi S, Higano S, Ishii K et al.: Hypoxic brain damage: cortical laminar necrosis and delayed changes in white matter at sequential MR-imaging. *Radiology* 1993; 189: 449–56
34. Ranaa L, Sooda D, Chauhanb R et al.: MR Imaging of hypoxic ischemic encephalopathy – Distribution Patterns and ADC value correlations. *European Journal of Radiology*; 2018; 5: 215–220
35. Mlynash M, Campbell DM, Leproust EM, et al.: Temporal and spatial profile of brain diffusion-weighted MRI after cardiac arrest. *Stroke* 2010; 41(8): 1665–72
36. Wijdicks EF, Campeau NG, Miller GM. MR imaging in comatose survivors of cardiac resuscitation. *AJNR Am J Neuroradiol* 2001;22:1561–1565.
37. Huang BY, Castillo M. Hypoxic-ischemic brain injury: imaging findings from birth to adulthood. *Radiographics*. 2008;28(2):417–39
38. Arbelaez A, Castillo M, Mukherji S: Diffusion-weighted MR imaging of global cerebral anoxia. *Am J Neuroradiol* 1999; 20: 999–1007
39. Sandroni C, D'Arrigo S, Cacciola S et al.: Prediction of poor neurological outcome in comatose survivors of cardiac arrest: a systematic review. *Intensive Care Med* 2020; 46: 1803–51
40. Lövblad KO, Wetzel SG, Somon T et al.: Diffusion-weighted MRI in cortical ischaemia. *Neuroradiology* 2004; 46: 175–82
41. Wu O, Sorensen AG, Benner T et al.: Comatose patients with cardiac arrest: Predicting clinical outcome with diffusion-weighted MR imaging. *Radiology*. 2009;252(1):173–81.
42. Wijman CAC, Mlynash M, Caulfield AF et al.: Prognostic value of brain diffusion- after weighted imaging cardiac arrest. *Annals of neurology* 2009; 65(4): 394–402
43. Hirsch KG, Mlynash M, Eyngorn I et al.: Multi-Center Study of Diffusion-Weighted Imaging in Coma After Cardiac Arrest. *Neurocrit Care* 2016; 24(1): 82–9
44. Moon HK, Jinhee J, Kyu, NP et al.: Quantitative analysis of relative volume of low apparent diffusion coefficient value can predict neurologic outcome after cardiac arrest. *Resuscitation* 2018; 126: 36–42
45. Hirsch KG, Fischbein N, Mlynash M, et al.: Prognostic value of diffusion-weighted MRI for post-cardiac arrest coma. *Neurology*. 2020 Apr 21; 94(16): e1684-e1692
46. Wouters A, Scheldeman L, Plessers S et al: Added Value of Quantitative Apparent Diffusion Coefficient Values for Neuroprognostication After Cardiac Arrest. *Neurology* 2021; 25; 96(21): e2611-e2618
47. Puybasset L, Perlberg V, Unrug J et al.: Prognostic value of global deep white matter DTI metrics for 1-year outcome prediction in ICU traumatic brain injury patients: an MRI-COMA and CENTER-TBI combined study. *Intensive Care Med* 2022; 48(2): 201–12
48. An C, You Y, Park JS et al.: The cut-off value of a qualitative brain diffusion-weighted image (DWI) scoring system to predict poor neurologic outcome in out-of-hospital cardiac arrest (OHCA) patients after target temperature management. *Resuscitation* 2020; 157: 202–10
49. Hirsch KG, Mlynash M, Jansen S et al.: Prognostic value of a qualitative brain MRI scoring system after cardiac arrest. *J Neuroimaging* 2015;25(3): 430–7
50. Wei R, Wang C, He F et al.: Prediction of poor outcome after hypoxic-ischemic brain injury by diffusion-weighted imaging: A systematic review and meta-analysis. *PLoS ONE* 2019; 14(12): e0226295
51. Cronberg T, Rundgren M, Westhall E et al.: Neuron-specific enolase correlates with other prognostic markers after cardiac arrest. *Neurology* 2011; 77(7): 623–30
52. Keijzer HM, Hoedemaekers CWE, Meijer FJA et al.: Brain imaging in comatose survivors of cardiac arrest: Pathophysiological correlates and prognostic properties. *Resuscitation* 2018; 133: 124–36
53. Sandroni C, D'Arrigo S, Cacciola S et al.: Prediction of good neurological outcome in comatose survivors of cardiac arrest: a systematic review. *Intensive Care Med* 2022 Mar 4. doi: 10.1007/s00134-022-06618-z. Online ahead of print
54. Dankiewicz J, Cronberg T, Lilja G et al.: Hypothermia versus Normothermia after Out-of-Hospital Cardiac Arrest. *N Eng J Med* 2021; 384: 2283–94